

THE ROLE OF APOPTOSIS IN PREECLAMPSIA

Levy R

**Department of Obstetrics and Gynecology, Kaplan Medical Center, affiliated to
the Hebrew University School of Medicine.**

Running title: Apoptotic processes are involved in the pathophysiology of
preeclampsia

Key words: Apoptosis, preeclampsia, placenta, hypoxia, p53.

Correspondence:

Roni Levy, MD, Department of Obstetrics and Gynecology, Kaplan Medical Center

Tel:08-9441558, 050-903984. Fax: 08-9467045. E mail: Roni_L@clalit.org.il

Preeclampsia occurs in 5-7% of pregnancies and is a major cause of perinatal and maternal morbidity and mortality worldwide. Preeclampsia is a condition specific to pregnancy presenting with a clinical picture of hypertension, proteinuria and different systemic manifestations. Despite intensive research the pathophysiology leading to the consequences of preeclampsia remain elusive. The most accepted concept is that immune maladaptation to the fetal allograft causes defective placentation. During preeclamptic pregnancy cytotrophoblast invasion of the myometrium is abnormally shallow, and remodeling of the uterine spiral artery is incomplete. Many vessels are occluded by atherosclerosis, with an accumulation of lipid-laden macrophages and a perivascular mononuclear cell infiltrate. The net result is uteroplacental ischemia.

A yet undefined toxic circulating factor released by the ischemic placenta enters the maternal circulation and damages vascular endothelium and is responsible for the clinical disease characterized by impaired endothelial function [1]. This circulating factor may be specific upregulated protein, cellular debris or free fetal DNA.

Numerous articles have been published addressing the association between altered apoptosis and preeclampsia [2-7]. Apoptosis has been linked to preeclampsia at different steps in the proposed pathway leading to the clinical manifestation of preeclampsia. At each of these steps, apoptosis may play a central role.

The term apoptosis, Greek for “falling off”—like autumn leaves—coined by Kerr, Wyllie and Currie in 1972 [8], is descriptive of a unique morphology of cell death. It is distinct from cell death by necrosis, where damage to groups of cells, cytoplasmic swelling and inflammatory processes are characteristic. Unlike necrosis, apoptosis includes chromatin condensation with nuclear fragmentation and cytoplasmic condensation with cell shrinkage. The morphological changes may be part of normal physiology or may be secondary to pathological insult. Apoptosis occurs during

normal embryonic development and in the turnover of mature tissues. However, apoptosis is also found in tissues exposed to exogenous stimuli, such as hypoxia or cytotoxic agents, and it can be induced by ligand-receptor interactions, such as with Fas and TNF- α [9]. Apoptosis is described in many physiological as well as pathological processes in the human body, including preeclampsia with or without fetal growth restriction (FGR). It may be argued that enhanced apoptosis may be a normal response to stimuli such as hypoxia. However several in-vitro and in-vivo studies showed that specific and unspecific scavengers of apoptosis such as caspase inhibitor and Epidermal growth factor decreased the level of apoptosis and improved the tissue function [10-12]. These studies point to an important role of apoptosis in pathologic processes.

In this article, I review the evidence linking apoptosis to each step in the mechanism of preeclampsia (Fig. 1).

Altered apoptosis induces maternal immune intolerance toward the fetus

The fetus and placenta are considered a semi-allograft to the mother's immune system and consequently generate an immune rejection response. From early pregnancy, there is a large influx of macrophages and lymphocytes into the decidua at the maternal-fetal interface such that white blood cells comprise up to 40% of the cells found in the decidua during pregnancy [13]. How is the developing fetus protected from these cells? One of the most accepted theories is that Human Leukocyte Antigen (HLA) -G has an immunologically permissive role of the antigenic mismatch between mother and fetus [14]. HLA-G expression was found to be defective in extravillous cytotrophoblast of preeclamptic placentas. Further, the serum levels of a soluble HLA-G1 isoform who downregulates T cell and NK activity by

inducing apoptosis, were decreased in preclamptic patients [14]. The exact mechanism by which HLA-G causes cell apoptosis is unknown.

Another theory that are taken from other systems of the body that are "immunologically preivileged sites". The anterior chamber of the eye and the testes are protected from antigens which penetrate these sites and fail to elicit an immune response. It turns out that cells in these sites differ from the other cells of the body in that they express high levels of FasL at all times. Thus antigen-reactive T cells, which express Fas, would be killed when they enter these sites[15]. Fas, a 45-kDa surface protein which is a member of the tumor necrosis factor superfamily, induces cell apoptosis upon binding to the ligand FasL [16]. Several pieces of evidence point to the key role played by the Fas-FasL system in protecting the fetus from maternal immune system [17]. The trophoblast of first-trimester placentas expresses FasL, while Fas antigen is localized mainly on decidual cells, in particular the maternal leukocytes [18,19]. When trophoblasts expressing FasL are exposed to activated lymphocytes in-vitro, they induce lymphocytes apoptosis [20]. In mice, FasL is positioned to prevent the exchange of activated immune cells between the mother and the fetus, and deters trafficking of activated Fas-expressing immune cells at the maternal-fetal interphase [21]. Mice lacking FasL (gld) exhibit leukocyte infiltration of the decidual-placental interface and increased fetal loss. In preeclampsia, decidual cells exhibit lower expression of Fas antigen [6], and lower expression of FasL is seen on trophoblast cells [4]. In addition, the level of serum-soluble Fas (sFas) antigen is elevated in the maternal circulation [6]. sFas, a spliced product of the Fas antigen, does not stimulate apoptosis; rather, it protects cells from apoptosis by competitive binding to the Fas antigen. Thus, elevated levels of sFas in the maternal circulation concomitant with lower expression of Fas and FasL result in diminished apoptotic

deletion of the lymphocytes in the maternal decidual system. As a consequence, there is increased apoptosis and trophoblast damage in the fetal placental compartment. Indeed, samples taken from the uterine wall of preeclamptic women demonstrated excess of macrophages. These macrophages when exposed to TNF- α , a known stimulator of placental apoptosis [22], may limit trophoblast invasion of spiral arterial segments by TNF- α mediate apoptosis [23]. Indeed TNF- α is found in high concentration of plasma, amniotic fluid and placental tissue of patients with preeclampsia [24,25]. The enhanced apoptosis of the invading trophoblasts results in limited invasion of the spiral arteries due to reduced activity of the cytotrophoblasts, so that myometrial segments of the spiral arteries remain intact resulting in the formation of an arteriolar system with high resistance. The failure of trophoblast invasion results in a reduction in uteroplacental perfusion, with the placenta becoming increasingly ischemic as gestation progresses. Placentas from women with preeclampsia display an increased frequency of placental infarcts and altered morphology, evidenced by abnormal cytotrophoblast proliferation and increased formation of syncytial knots [26]. In a study [27] comparing tissue samples of the villus-uterus attachment, taken from preeclamptic and control patients, samples from the former showed widespread apoptosis, while those from normal pregnancies had a low rate of apoptotic cells. Further, the samples taken from preeclamptic patients showed lower expression of one of the proapoptotic proteins, Bcl-2. Increased apoptosis along with decreased expression of Bcl-2 has also been reported in myocardial cells [12]. Thus, excess apoptotic activity in placental bed of preeclamptic women inhibits trophoblast invasion into the spiral artery by increasing trophoblast apoptosis.

Several mechanisms, in addition to apoptosis, have been reported to limit extravillous invasion of the placental bed. These consist of the reduced expression of various proteins, among them integrin $\alpha 1/\beta 1$, matrix metalloproteinase, vascular cell adhesion molecules, Vascular endothelial growth factor (VEGF) and heparin-binding Epidermal growth factor (EGF) [28]. A reduction in the expression of these proteins necessary for extravillous trophoblast differentiation along the invasive pathway is compatible with the view that cells entering the apoptotic cascade downregulate their levels of protein transcription.

Hypoxia induces placental apoptosis

Placental ischemia secondary to defective placentation may be a prerequisite for the development of preeclampsia. Human term trophoblasts exposed to hypoxia in vitro exhibit enhanced apoptosis [11]. The process was associated with increased expression of the proapoptotic proteins p53 and Bax and reduced expression of the antiapoptotic Bcl-2. Enhanced apoptosis and upregulation of p53 were also found in placental samples taken from pregnancies complicated by preeclampsia [29]. Hypoxia triggers apoptosis via a mechanism that involves predominantly mitochondrial pathways, as opposed to ligand-receptor pathways which are mediated by cytokines such as TNF- α or Fas ligand (Fig 1). In this pathway, death signals are stimulated via modulation of the expression of specific apoptotic-related genes such as p53 and proteins of the Bcl-2 family. p53 plays a pivotal role in the cellular response to DNA damage, halting the cell cycle to allow repair of DNA. If repair is not possible, p53 promotes apoptosis. p53 is an unstable protein with a short half-life: exogenous stimuli such as hypoxia and oxidative stress stabilize it . p53 plays an important role in hypoxia-induced cell death in diverse cell types, including cardiocytes, hepatocytes

and neuronal cells [30]. p53 expression is enhanced in cultured trophoblasts exposed to hypoxia [11] as well as in placental biopsies taken from pregnancies complicated by preeclampsia and FGR.

Hypoxia may not be the sole inducer of placental apoptosis. Oxidative free radicals are frequently, although not universally, increased in preeclampsia [1]. Retention of vasoreactivity of the spiral arteries caused by defective placentation may result in intermittent perfusion of the intervillous space, fluctuating oxygen tension, and ischemia-reperfusion insult of the villus. This oxidative stress may be associated with enhanced placental apoptosis and increased turnover of the syncytiotrophoblast [31].

Thromboxane A2 levels are elevated in the circulation of pregnant preeclamptic women and in the placental villi [32]. Elevated levels of thromboxane A2 are associated with thrombocyte aggregation, which may create a predisposition to enhanced thrombosis and infarct injury of the placenta. Thromboxane A2 has been recently found to enhance apoptosis in primary-term human trophoblasts [33].

Placental apoptosis increases syncytiotrophoblast deportation

One of the unsolved questions in the pathogenesis of preeclampsia is the link between placental ischemia and endothelial cell dysfunction. Systemic endothelial damage appears to be the central theme in the signs and symptoms of preeclampsia. One of the theories, presented by Redman and Sargent [34], is that systemic endothelial damage is caused by microdeposition of syncytiotrophoblast microvillous membrane particles. These particles can be detected in the plasma of normal pregnancies but increase in women with preeclampsia. The increased syncytiotrophoblast deportation in preeclampsia may be caused by enhanced apoptosis at the syncytium affecting the integrity of the tissue. It has been proposed

that apoptosis plays an important role in the renewal of the syncytium [35,36]. Apoptotic nuclei are found in syncytial knots and probably contribute to the shedding of syncytial fragments into the maternal circulation [37]. This process is enhanced in the syncytium of placentas from pregnancies complicated by preeclampsia [6,29]. When the maternal circulation cannot compensate for this enhancement, systemic endothelial damage occurs.

Altered apoptosis in preeclampsia is found not only in the placenta but also in the endothelial cells [38]. Endothelial-tissue apoptosis may occur from the toxic effect of excessive syncytial fragments, but it may also occur from the local hypoxic environment caused by the constriction of efferent blood vessels or from the effect mediated by free radicals. This effect may occur via p53-mediated endothelial apoptosis, as found in other systems (11,12) . Another hypothesis is that increased secretion of TNF- α induces activation and apoptosis of endothelial tissue [39]. Another hint for the apoptosis induced trophoblast microfragments deportation is the increased level of free DNA in the circulation of patient with preeclampsia [40].

In summary, altered apoptosis is involved in each step of the pathogenesis of preeclampsia. While deficient apoptosis may induce a maternal immune response against the fetus, enhanced apoptosis may interfere with the process of placentation, placental ischemia and subsequently, systemic endothelial damage. Thus, treatment modalities to inhibit or accelerate apoptosis cannot be employed in early pregnancy for prevention. Later in pregnancy, it would make sense to test treatments modalities that inhibit hypoxia mediated apoptosis in patients who early testing, such as abnormal Doppler of the uterine artery, indicates the initiation of preeclampsia. This however has to be first tested in animals models.

Since multiple different mechanisms, rather than a single factor, could contribute to the development of apoptosis, further studies to clarify the signaling mechanisms of apoptosis in preeclampsia should be conducted before any investigational treatment modalities are employed.

References

1. Dekker GA, Sibai BM. Etiology and pathogenesis of preeclampsia: current concepts. *Am J Obstet Gynecol* 1998; 179: 1359-75.
2. Leung DN, Smith SC, To KF, Sahota DS, Baker PN. Increased placental apoptosis in pregnancies complicated by preeclampsia. *Am J Obstet Gynecol* 2001; 184 (96): 1249-50.
3. Harirah H, Donia SE, Hsu CD. Serum soluble Fas in the syndrome of hemolysis, elevated liver enzymes, low platelets. *Obstet Gynecol* 2001; 98(2): 295-8.
4. Allaire AD, Ballenger KA, Wells SR, McMahon MJ, Lessey BA. Placental apoptosis in preeclampsia. *Obstet Gynecol* 2000; 96: 271-6.
5. Hsu CD, Harirah H, Basherra H, Mor G. Serum soluble Fas levels in preeclampsia. *Obstet Gynecol* 2001; 97(4): 530-2.
6. Ishihara N, Matsuo H, Murakoshi H, Laoag-Fernandez JB, Samoto T, Maruo T. Increased apoptosis in the syncytiotrophoblast in human term placentas complicated by either preeclampsia or intrauterine growth retardation. *Am J Obstet Gynecol* 2002; 186(1): 158-66.
7. Darmochwal-Kolarz D, Rolinski J, Leszczynska-Gorzela B, Oleszczuk J. Fas antigen expression on the decidual lymphocytes of pre-eclamptic patients. *Am J Reprod Immunol* 2000; 43(4): 197-201.
8. Kerr JFR, Wylie AH, Currie AR. Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer* 1972; 26: 239-57.
9. Green DR. Apoptotic pathways: the roads to ruin. *Cell* 1998; 94: 695-8.

10. Levy R, Smith DS, Chandler K, Sadovsky Y, Nelson DM. Apoptosis in human cultured trophoblasts is enhanced by hypoxia and diminished by epidermal growth factor. *Am J Physiol Cell Physiol* 2000; 278: C982-8.
11. Banasiak KJ, Haddad GG. Hypoxia-induced apoptosis: effect of hypoxic severity and role of p53 in neuronal cell death. *Brain Res* 1998;797:295-304
12. Long X, Boluyt A, Hipolito B, Lundberg M, Zheng J, Oíniel L, et al. P53 and the hypoxia-induced apoptosis of cultured neonatal rat cardiac myocytes. *J Clin Invest* 1997;99:2635-43.
13. Bulmar J, Johnson P. Immunohistological characterization of the decidual leukocytes infiltrate related to endometrial gland epithelium in early human pregnancy. *Immunology* 1985; 55: 35-44.
14. Le Bouteiller P, Pizzato N, Barakonyi A, Solier C. HLA-G, preeclampsia, immunity and vascular events. *J of Reprod. Immun.* 2003; 59: 219-34.
15. Nagata S. Apoptosis by death factor. *Cell* 1997; 88: 355-65.
16. Griffith TS, Brunner T, Fletcher SM, Green DR, Ferguson TA. Fas ligand-induced apoptosis as a mechanism of immune privilege. *Science* 1995; 270(5239): 1189-92.
17. Pearson H, Reproductive immunology: Immunity's pregnant pause. *Nature*. 2002; 21(420) 6913: 265-6.
18. Runic R, Lockwood CJ, Ma Y, Dipasquala B, Guller S. Expression of Fas ligand by human cytotrophoblasts: implications in placentation and fetal surviving. *J Clin Endocrinol Metab* 1996; 81: 3119-22.

19. Uckan D, Steele A, Wang B-Y, Chamizo W, Koutsonikolis A, Gilbert-Barness E, Good RA. Trophoblasts express Fas ligand: a proposed mechanism of immune privilege in placenta and maternal invasion. *Mol Hum Reprod* 1997; 3: 655-62.
20. Hammer A, Blaschitz A, Daxböck C, Walcher W, Dohr G. Fas and Fas-ligand are expressed in the uteroplacental unit of first trimester pregnancy. *Am J Reprod Immunol* 1999; 41: 41-51.
21. Hunt JS, Vassmer D, Ferguson TA, Miller L. Fas ligand is positioned in mouse uterus and placenta to prevent trafficking of activated leukocytes between the mother and the conceptus. *J Immunol* 1997; 158: 4122-8.
22. Yui J, Garcia-Lloret M, Wegmann TG, Guilbert LJ. Cytotoxicity of tumour necrosis factor-alpha and gamma-interferon against primary human placental trophoblasts. *Placenta* 1994;15:819-35.
23. Reister F, Frank H-G, Kingdom JCP, Heyl W, Kaufman P, Rath W, Huppertz B. Macrophage-induced apoptosis limits endovascular trophoblast invasion in the uterine wall of preeclamptic women. *Laboratory Investigation* 2001; 81:1143-52.
24. Wang Y, Walsh SW. TNF α concentrations and mRNA expression are increased in preeclamptic placentas. *J Reprod Immunol* 1996;32:157-69.
25. Fisher SJ, Roberts JM. Defects in placentation and placental perfusion. In Linheimer MD, Roberts JM, Cunningham FG. *Chesley's hypertensive disorders in pregnancy*. 2nd ed. Stanford, CT: Appleton & Lange, 1999; 377-94.

26. Nelson DM, Johnson RD, Smith SD, Anteby EY, Sadovsky Y.
Hypoxia limits differentiation and upregulates expression and activity of PGH synthase 2 in culture trophoblast from term human placenta.
Am J Obstet Gynecol 1999; 180: 896-902.
27. DiFederico E, Genbacev O, Fisher SJ. Preeclampsia is associated with widespread apoptosis of placental cytotrophoblasts within the uterine wall. Am J Pathol 1999; 155(1): 293-7.
28. Leach RE, Romero R, Kim YM, Chaiworapogsa T, Kilburn B, Dey SK, Johnson A, Armant DR. Preeclampsia and expression of heparin-binding EGF like growth factor. Lancet 2002; 1936(9341): 1215-9.
29. Levy R, Smith DS, Sadovsky Y, Nelson DM. Apoptosis and p53 are enhanced in placentas from pregnancies complicated by fetal growth restriction induced by hypoxia. Am J Obstet Gynecol 2002; 186(5): 1056-61.
30. Stempian-Oteero A, Karsan A, Cornejo CJ, Xiang H, Eunson T, Morrison RS et al. Mechanism of hypoxia-induced endothelial cell death. J Cell Biol 1999; 274: 8039-45.
31. Crocker IP, Cooper S, Ong SC, Baker PN. Differences in apoptotic susceptibility of cytotrophoblasts and syncytiotrophoblast in normal pregnancy to those complicated by preeclampsia and intrauterine growth restriction. Am J Pathol 2003; 162 (2) :637-43.
32. Roberts JM, Taylor RN, Goldfien A. Clinical and biochemical evidence of endothelial cell dysfunction in the pregnancy syndrome preeclampsia. Am J Hypertension 1991; 4: 700-8.

33. Kamran Y, Smith DS, Levy R, Schaiff WT, Sadovsky Y, Nelson DM. Thromboxane A2 limits differentiation and enhances apoptosis of cultured human trophoblast. *Pediatr Res* 2001; 50(2): 203-9.
34. Redman CW, Sargent IL. Placental debris, oxidative stress and preeclampsia. *Placenta* 2000; 21(7): 597-602.
35. Nelson DM. Apoptotic changes occur in the syncytiotrophoblast of human placental villi where fibrin type fibrinoid is deposited at discontinuities in the villous trophoblast. *Placenta* 1996; 17: 387-91.
36. Huppertz B, Frank HG, Kingdom JCP, Reister F, Kaufman P. Villous cytotrophoblast regulation of the syncytial apoptotic cascade in the human placenta. *Histochem Cell Biol* 1998; 110: 495-508.
37. Mayhew TM, Leach L, McGee R, Ismail WW, Myklebust R, Lammiman MJ. Proliferation, differentiation and apoptosis in villous trophoblast at 13-41 weeks of gestation. *Placenta* 1999; 20: 407-2.
38. Neale D, Demasio K, Illuzi J, Chaiworapongsa T, Romero R, Mor G. Maternal serum of women with pre-eclampsia reduces trophoblast cell viability: evidence for an increased sensitivity to Fas-mediated apoptosis. *J Matern Fetal Neonatal Med* 2003;13(1): 39-44.
39. Hung TH, Charnock-Jones DS, Skepper JN, Burton GI. Secretion of tumor necrosis factor-alpha from human placental tissues induced by hypoxia-reoxygenation causes endothelial cell activation in vitro: a potential mediator of the inflammatory response in preeclampsia. *Am J Pathol* 2004; 164(3):1049-61.
40. Levine RJ, Qian C, Leshane Es, Yu KF, England LJ, Schisterman EF, Wataganara T, Romero R, Bianchi DW. Two stage elevation of cell-

free fetal DNA in maternal sera before onset of preeclampsia. Am J Obstet Gynecol 2004; 190 (3):707-13.

Figure 1. Apoptotic pathways. There are two main apoptotic pathways. One induced by internal or external stimuli mediated by proteins from the Bcl-2 family, via the mitochondria, the other through ligand to receptor stimuli such as Fas and TNF- α .




Figure 2. Apoptosis and sequences of preeclampsia.

