General examination

| | Weight (kg) | Abdomen circumference (sm) | Height of fundus (sm) | Blood pressure | | Heat rate | Edema | Proteinuria | Gestosis |
|----------|----------------|----------------------------------|-----------------------------|-------------------------------------|--|-------------------------|----------|-------------|----------|
| rameters | | | | R. hand Sist/diast and points | L. hand Sist/diast And points | (impacts one minute) | (points) | (points) | Index |
| Pa | | | | | 5746 | 100 | 1425 | | |

Ryodoraku

| R TE | Har | nds | | Legs | |
|----------------------|-------|------|-----------------------|-------|------|
| Channels | Right | Left | Channels | Right | Left |
| P ₉ (LU) | | | RP ₃ (SP) | | |
| MC ₇ (P) | | | F ₃ (LIV) | | |
| C ₇ (HE) | | | R ₃ (KID) | | |
| IG ₄ (SI) | | | V ₆₅ (BL) | | |
| TR ₄ (SJ) | | | VB ₄₀ (GB) | | |
| GI ₅ (LI) | | | E ₄₂ (ST) | | |

The examination is repeated 2-3 times till deliver. We recommend do it during first visit, on 18-19th, 32-33rd weeks and just before labor. It is necessary to mark the rise of weight for all period of pregnancy.